

Flex Therapist CEUs

Changing Landscape of Outpatient Physical Therapy

1. Traditional care models are associated with all of the following negative healthcare outcomes EXCEPT:

- A. Excessive medical costs
 - B. Increased frequency of high risk surgeries
 - C. Decreased episodes of care
 - D. Poor access to physical therapy services
-

2. The Affordable Care Act and Comprehensive Care for Joint Replacement programs encourage the use of which type of care model?

- A. Value-based
 - B. Traditional
 - C. Empirically-driven
 - D. Primary
-

3. Rising healthcare expenditures can be attributed to which of the following factors:

- A. Specialty practice referrals
 - B. Steroid injections
 - C. Prescription of opioid medications
 - D. All of the above
-

4. Which of the following factors would be considered a disruption to traditional care models?

- A. Physician-based referrals to physical therapy
 - B. Early physical therapy models
 - C. Physician-based referrals to specialists
 - D. Early mental health interventions
-

5. Recent clinical practice guidelines for non-pharmacological treatments of low back pain have contributed to the evolution of outpatient services by affecting which of the following components of therapy:

- A. Shifting the timing of interventions

- B. Rearranging the order of services**
 - C. Altering access routes**
 - D. All of the above**
-

6. Changes to referral patterns have dictated new patient care responsibilities to clinicians in the outpatient setting. Which of the following scenarios is most reflective of this transition?

- A. COVID-19 has forced many patients to seek outpatient physical therapy services as a last resort to medical care.**
 - B. Increased acuity of patient referrals requires therapists to be knowledgeable about wound care management.**
 - C. The Affordable Care Act altered the timing of physical therapy referrals and prioritizes physician referrals.**
 - D. Bundled payments have increased the number of patients who are being sent to post-acute facilities.**
-

7. Clinicians should perform a review of systems to identify conditions that may not be amenable to therapeutic interventions. Which of the following questions would be an appropriate review of systems for a patient who presents with low back pain?

- A. Have you recently experienced excessive thirst?**
 - B. Have you recently experienced lightheadedness?**
 - C. Have you recently experienced night pain?**
 - D. Have you recently experienced changes in skin texture?**
-

8. Under the clinical practice guideline, clinician role in the presence of a potential lower extremity deep venous thrombosis (DVT) include all of the following EXCEPT:

- A. Alert the physician for the need to perform a thorough examination**
 - B. Perform screening measures for fall risk**
 - C. Provide patient education regarding risk factors for developing a DVT**
 - D. Recommend use of anticoagulants to achieve therapeutic threshold levels**
-

9. Which of the following factors is considered to be indicative of a systemic infection from inadequate wound healing?

- A. 100 degree fever**
 - B. Diarrhea**
 - C. Odorous incision site**
 - D. Swollen and itchy wound**
-

10. A review of symptoms with the main objective of identifying potential non-

musculoskeletal system involvement is known as:

- A. Red flag screening**
 - B. Tests and measures**
 - C. Yellow flag screening**
 - D. Standardized measures**
-

11. Which of the following goals accurately describes the objective of the Clinical Practice Guideline for Venous Thromboembolism/Lower Extremity Deep Vein Thrombosis?

- A. Enables clinicians to identify the presence of Postthrombotic Syndrome**
 - B. Encourages clinicians to participate in an interdisciplinary approach for safe mobility**
 - C. Informs clinicians on ways to identify when the patient's therapeutic threshold levels are not met**
 - D. All of the above**
-

12. Which of the following scenarios reflect the importance of knowing how incisional closures affect wound management?

- A. Erythema after staple removal is abnormal and should be considered a localized infection.**
 - B. Steri-strips should be removed by a medical professional after 7-10 days following surgery.**
 - C. Tissue adhesives offer the strongest protection against foreign bacteria and infection.**
 - D. Patients with risk factors for poor wound healing will most likely have a staple closure.**
-

13. Screening tools to assist clinicians when administering screenings for depressive symptoms include:

- A. Beck Depression Inventory**
 - B. Center for Epidemiologic Studies Depression Scale**
 - C. Hamilton Depression Rating Scale**
 - D. All of the above**
-

14. Under the Merit-based Incentive Payment System, physical therapists who report quality measures for depression screening must include which of the following documentation items:

- A. Suicide Risk Assessment**
- B. Pharmacological interventions**
- C. Treatment for depression**

D. All of the above

15. Performing a medical reconciliation can identify negative behaviors regarding medication usage including:

- A. Polypharmacy**
 - B. Non-adherence**
 - C. Adverse drug events**
 - D. All of the above**
-

16. Which of the following patient descriptors increases the risk of poor adherence to medication regime?

- A. Patients who live with an elderly spouse**
 - B. Patients who take less than 2 medications per day**
 - C. Patients who are diagnosed with clinical depression**
 - D. Patients who use two or more pharmacies to fulfill prescriptions**
-

17. Clinicians should prioritize medication reconciliation in order to:

- A. Maintain patient safety when under the care of a physical therapist or physical therapy assistant**
 - B. Obtain the correct information regarding patient's responses to prescribed pharmaceuticals**
 - C. Correct physician error when prescribing an incorrect dosage**
 - D. Confirm that the patient is utilizing one pharmacy to fulfill prescription drugs, especially in the presence of polypharmacy**
-

18. Which of the following statements best describes the importance of understanding an expected physiological response to exercise?

- A. Clinicians can detect adverse drug reactions that may cause exercise-induced changes.**
 - B. Clinicians can identify limitations to aerobic activity that can serve as a foundation for exercise prescription.**
 - C. Clinicians can screen for underlying pathologies that may affect a patient's capacity for exercise.**
 - D. Clinicians can modify exercise parameters to encourage exercise-induced fatigue.**
-

19. Physical therapy roles during disaster mitigation may include:

- A. Assisting with emergency amputations**
- B. Administering tourniquettes to wounds**

- C. Prescribing anti-inflammatory medications**
 - D. Triaging patients who require surgery**
-

20. A 60 year old man is exercising on the treadmill with a resting heart rate of 75 beats per minute (BPM). What would be an appropriate heart rate response to this exercise?

- A. Heart rate should rise incrementally by 15 BPM**
 - B. Heart rate should decrease to resting level within 15 minutes following exercise**
 - C. Heart rate should rise incrementally by 10 BPM**
 - D. Heart rate should decrease to resting level within 10 minutes following exercise**
-

Copyright © 2024 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>