

# Flex Therapist CEUs

## Management of ACL Injuries in Children

**1. How many main ligaments surround the knee to provide support?**

- A. 4
  - B. 1
  - C. 3
  - D. 2
- 

**2. What is not a risk factor for ACL tears?**

- A. Improper footwear
  - B. Female gender
  - C. Sports
  - D. Hip weakness
- 

**3. A major immediate sign of an ACL tear is;**

- A. Nausea
  - B. Falling
  - C. A popping sound
  - D. Burning pain
- 

**4. What is not a test used to determine a torn ACL?**

- A. Reverse Clunk Test
  - B. Reverse Pivot Shift Test
  - C. Lachman's Sign Test
  - D. Clunk Test
- 

**5. What can be used to show the extent of injury to the surrounding cartilage?**

- A. X-ray
  - B. MRI
  - C. Anterior Drawer Test
  - D. CT scan
- 

**6. What method is useful for alleviating pain and swelling throughout the process?**

- A. SALT

- B. TIME
  - C. POLICE
  - D. FORT
- 

**7. What movement is unlikely to contribute to an ACL injury?**

- A. Pivoting
  - B. Cutting
  - C. Jumping
  - D. Squatting
- 

**8. How long does it normally take to recover from ACL reconstruction?**

- A. 3-4 months
  - B. 6-12 months
  - C. 6-24 months
  - D. 8-10 months
- 

**9. Disturbance to the growth plate can result in leg length discrepancies and angular deformities.**

- A. True
  - B. False
- 

**10. Immediately after surgical reconstruction, the patient should begin exercise with resistance.**

- A. True
  - B. False
- 

Copyright © 2025 Flex Therapist CEUs

Visit us at <https://www.flextherapistceus.com>